arXiv:quant-ph/0405176v1 28 May 2004

The Evolution Operator of The Three Atoms Tavis-Cummings Model

Jin-fang Cai

Department of Applied Physics, Beijing Institute of Technology,
Beijing, 100081, P.R. China

Abstract

The explicit form of evolution operator of the three atoms Tavis-Cummings Model is given.

The Tavis-Cummings Models have been widely studied in the field of quantum optic and quantum computer. In this brief letter, The analytical form of the evolution operators of T-C model with three atoms is given. Compared with the result in the paper[1], our result is more convenient for further application.

Under the rotating-wave approximation and resonant condition, the Hamiltonian of the Tavis-Cummings model with three atoms in the interaction picture is

$$\hat{H}_I = \hbar\gamma \sum_{i=1}^{3} (\hat{a}\hat{\sigma}_i^{+} + \hat{a}^{\dagger}\hat{\sigma}_i^{-}) \tag{1}$$

We denote the ground and exited states for the atom by $|g\rangle$ and $|e\rangle$. The evolution operator $\hat{U}(t) = \exp(-i\hat{H}t/\hbar)$ in the atomic basis $|eee\rangle, |eeg\rangle, |ege\rangle, |gee\rangle, |egg\rangle, |geg\rangle, |gge\rangle, |ggg\rangle$, is found to be

$$\begin{pmatrix} \hat{U}_{11} & \cdots & \hat{U}_{18} \\ \vdots & \ddots & \vdots \\ \hat{U}_{81} & \cdots & \hat{U}_{88} \end{pmatrix} \tag{2}$$

where

$$\begin{aligned}
\hat{U}_{11} &= \frac{(7+2\hat{N}+\hat{\Omega})\cos(\hat{\Theta}_1\gamma t) + (-7-2\hat{N}+\hat{\Omega})\cos(\hat{\Theta}_2\gamma t)}{2\hat{\Omega}} \equiv u_{11}(\hat{N}) \\
\hat{U}_{22} &= \hat{a}^{\dagger}\frac{\left(-1-2\hat{N}+\hat{\Omega}\right)\cos(\hat{\Theta}_1\gamma t) + \left(1+2\hat{N}+\hat{\Omega}\right)\cos(\hat{\Theta}_2\gamma t) + 4\,\hat{\Omega}\,\cos(\sqrt{2+\hat{N}}\,\gamma t)}{6\,\hat{\Omega}(\hat{N}+1)}\hat{a} \\
&\equiv \hat{a}^{\dagger}\frac{u_{22}(\hat{N})}{\hat{N}+1}\hat{a} \\
\hat{U}_{88} &= \hat{a}^{\dagger 3}\frac{\left(1+2\hat{N}+\hat{\Omega}\right)\cos(\hat{\Theta}_1\gamma t) + \left(-1-2\hat{N}+\hat{\Omega}\right)\cos(\hat{\Theta}_2\gamma t)}{2\,\hat{\Omega}(\hat{N}+1)(\hat{N}+2)(\hat{N}+3)}\hat{a}^3
\end{aligned}$$

$$
\begin{aligned}
&\equiv \hat{a}^{\dagger 3}\frac{u_{88}(\hat{N})}{(\hat{N}+1)(\hat{N}+2)(\hat{N}+3)}\hat{a}^3\\
\hat{U}_{12} &= \frac{\hat{\Theta}_1\left(7+2\hat{N}+\hat{\Omega}\right)\sin(\hat{\Theta}_1\gamma t)+\hat{\Theta}_2\left(-7-2\hat{N}+\hat{\Omega}\right)\sin(\hat{\Theta}_2\gamma t)}{6i(1+\hat{N})\,\hat{\Omega}}\hat{a} \equiv \frac{u_{12}(\hat{N})}{\sqrt{\hat{N}+1}}\hat{a}\\
\hat{U}_{15} &= \frac{-\cos(\hat{\Theta}_1\gamma t)+\cos(\hat{\Theta}_2\gamma t)}{\hat{\Omega}}\hat{a}^2 \equiv \frac{u_{15}(\hat{N})}{\sqrt{(\hat{N}+1)(\hat{N}+2)}}\hat{a}^2\\
\hat{U}_{25} &= \hat{a}^{\dagger}\frac{-\hat{\Theta}_1\,(2+\hat{N})\sin(\hat{\Theta}_1\gamma t)+\hat{\Theta}_2\,(2+\hat{N})\,\sin(\hat{\Theta}_2\gamma t)+\sqrt{2+\hat{N}}\,\hat{\Omega}\,\sin(\sqrt{2+\hat{N}}\,\gamma t)}{3i(1+\hat{N})(2+\hat{N})\,\hat{\Omega}}\hat{a}^2\\
&\equiv \hat{a}^{\dagger}\frac{u_{25}(\hat{N})}{(\hat{N}+1)\sqrt{\hat{N}+2}}\hat{a}^2\\
\hat{U}_{58} &= \hat{a}^{\dagger 2}\frac{\hat{\Theta}_1\left(1+2\hat{N}+\hat{\Omega}\right)\sin(\hat{\Theta}_1\gamma t)+\hat{\Theta}_2\left(-1-2\hat{N}+\hat{\Omega}\right)\sin(\hat{\Theta}_2\gamma t)}{6i(1+\hat{N})(2+\hat{N})(3+\hat{N})\,\hat{\Omega}}\hat{a}^3\\
&\equiv \hat{a}^{\dagger 2}\frac{u_{58}(\hat{N})}{(\hat{N}+1)(\hat{N}+2)\sqrt{\hat{N}+3}}\hat{a}^3\\
\hat{U}_{18} &= \frac{i\hat{\Theta}_1\hat{\Theta}_2\left(\hat{\Theta}_2\sin(\hat{\Theta}_1\gamma t)-\hat{\Theta}_1\sin(\hat{\Theta}_2\gamma t)\right)}{3(\hat{N}+1)(\hat{N}+3)\,\hat{\Omega}}\hat{a}^3 \equiv \frac{u_{18}(\hat{N})}{\sqrt{(\hat{N}+1)(\hat{N}+2)(\hat{N}+3)}}\hat{a}^3
\end{aligned}
$$

and

$$
\begin{aligned}
&\hat{U}_{33}=\hat{U}_{44}=\hat{U}_{22};\\
&\hat{U}_{55}=\hat{U}_{66}=\hat{U}_{77}=\hat{a}^{\dagger 2}\frac{u_{22}(\hat{N})-\frac{1}{\hat{\Omega}}(\cos(\hat{\Theta}_1\gamma t)-\cos(\hat{\Theta}_2\gamma t))}{(\hat{N}+1)(\hat{N}+2)}\hat{a}^2\\
&\hat{U}_{23}=\hat{U}_{24}=\hat{U}_{32}=\hat{U}_{34}=\hat{U}_{42}=\hat{U}_{43}=\hat{a}^{\dagger}\frac{u_{22}(\hat{N})-\cos(\sqrt{\hat{N}+2}\gamma t)}{\hat{N}+1}\hat{a}\\
&\hat{U}_{56}=\hat{U}_{57}=\hat{U}_{65}=\hat{U}_{67}=\hat{U}_{75}=\hat{U}_{76}=\hat{a}^{\dagger 2}\frac{u_{22}(\hat{N})-\frac{1}{\hat{\Omega}}(\cos(\hat{\Theta}_1\gamma t)-\cos(\hat{\Theta}_2\gamma t))-\cos(\sqrt{2+\hat{N}}\gamma t)}{(\hat{N}+1)(\hat{N}+2)}\hat{a}^2\\
&\hat{U}_{13}=\hat{U}_{14}=\hat{U}_{12}; \qquad \hat{U}_{21}=\hat{U}_{31}=\hat{U}_{41}=\hat{a}^{\dagger}\frac{u_{12}(\hat{N})}{\sqrt{\hat{N}+1}}\\
&\hat{U}_{15}=\hat{U}_{16}=\hat{U}_{17}; \qquad \hat{U}_{51}=\hat{U}_{61}=\hat{U}_{71}=\hat{a}^{\dagger 2}\frac{u_{15}(\hat{N})}{\sqrt{(\hat{N}+1)(\hat{N}+2)}}\\
&\hat{U}_{28}=\hat{U}_{38}=\hat{U}_{48}=\hat{a}^{\dagger}\frac{u_{15}(\hat{N})}{(\hat{N}+1)\sqrt{(\hat{N}+1)(\hat{N}+2)}}\hat{a}^3;\\
&\hat{U}_{82}=\hat{U}_{83}=\hat{U}_{84}=\hat{a}^{\dagger 3}\frac{u_{15}(\hat{N})}{(\hat{N}+1)\sqrt{(\hat{N}+1)(\hat{N}+2)}}\hat{a}\\
&\hat{U}_{25}=\hat{U}_{26}=\hat{U}_{35}=\hat{U}_{37}=\hat{U}_{46}=\hat{U}_{47};
\end{aligned}
$$

$$\hat{U}_{52} = \hat{U}_{53} = \hat{U}_{62} = \hat{U}_{64} = \hat{U}_{73} = \hat{U}_{74} = \hat{a}^{\dagger 2}\frac{u_{25}(\hat{N})}{(\hat{N}+1)\sqrt{\hat{N}+2}}\hat{a}$$

$$\hat{U}_{27} = \hat{U}_{36} = \hat{U}_{45} = \hat{a}^{\dagger}\frac{u_{25}(\hat{N}) + \sin(\sqrt{2+\hat{N}}\gamma t)}{(1+\hat{N})\sqrt{2+\hat{N}}}\hat{a}^2;$$

$$\hat{U}_{54} = \hat{U}_{63} = \hat{U}_{72} = \hat{a}^{\dagger 2}\frac{u_{25}(\hat{N}) + \sin(\sqrt{2+\hat{N}}\gamma t)}{(1+\hat{N})\sqrt{2+\hat{N}}}\hat{a}$$

$$\hat{U}_{68} = \hat{U}_{78} = \hat{U}_{58}; \qquad \hat{U}_{85} = \hat{U}_{86} = \hat{U}_{87} = \hat{a}^{\dagger 3}\frac{u_{58}(\hat{N})}{(\hat{N}+1)(\hat{N}+2)\sqrt{\hat{N}+3}}\hat{a}^2$$

$$\hat{U}_{81} = \hat{a}^{\dagger 3}\frac{u_{18}(\hat{N})}{\sqrt{(\hat{N}+1)(\hat{N}+2)(\hat{N}+3)}}$$

where $\hat{\Omega}, \hat{\Theta_1}, \hat{\Theta_2}$ are defined by

$$\hat{\Theta_1}^2 = 5(\hat{N}+2) - \hat{\Omega} \qquad \hat{\Theta_2}^2 = 5(\hat{N}+2) + \hat{\Omega}$$

$$\hat{\Omega}^2 = 9 + 16(\hat{N}+2)^2$$

and $\hat{N} = \hat{a}^{\dagger}\hat{a}$

References

[1] K. Fujii, K. Higashida, R. Kato, T. Suzuki and Y. Wada, ”Explicit Form of Evolution Operator of Three Atoms Tavis-Cummings Model”, quant-ph/0404034